

's Crust of Bulgaria by P Receiver Functions

Receiver function technique is applied to evaluate the Moho depth and V_p/V_s ratio on the territory of Bulgaria. Bulgarian National Seismological Network (BNSN) is equipped with broad-band seismometers and digital acquisition systems. It enables application of modern techniques of analysis of the velocity structure in Bulgaria. The method was applied to eleven stations of BNSN.

The obtained results show that the Moho depth varies between 30 and 50 km. The crust is shallower in North-east of Bulgaria and goes deeper in Southwest direction. The crust is thicker in the Rhodopes massif, where it reaches more than 50 km beneath the station Musomishte (MMB). A thick sedimentary layer was also delineated in the Northern Bulgaria. The crustal structure in Southwestern Bulgaria especially beneath the station Vitosha (VTS) is very complex. Further more detailed analysis of data should be performed in order to better estimate the depth of the Mohorovicic discontinuity. The V_p/V_s ratio in the study area varies between 1.60 and 1.90. It varies between 1.80 and 1.90 in Northern Bulgaria and decreases in South and Southwest direction. The western part of Rhodopes massif is characterized by low V_p/V_s (1.6-1.65) and could be explained with presence of felsic rock formation.

Primary author: GEORGIEVA, Gergana (Sofia University St. Kliment Ohridski)

Presenter: GEORGIEVA, Gergana (Sofia University St. Kliment Ohridski)

Track Classification: Theme 1: The Earth as a Complex System