



ID:

Type: **Oral**

the Reduced Displacement Potentials of DPRK Nuclear Explosions Using Waveform Equalization Technique

Friday, 28 June 2019 12:00 (15 minutes)

We equalize regional P, surface, and the P+surface waves simultaneously to estimate reduced displacement potential (RDP) and depth of burial (DOB) of DPRK nuclear explosions, using waveforms from the stations at IRIS DMC. RDPs are predicted using the formula in Saikia (2017). The algorithm starts with the initial RDP and DOBs for two explosions. RDP of one explosion S_1 is convolved with the other explosion recorded data O_2 , and vice-versa (i.e., S_2 with O_1), which generates two convolution seismograms: S_1O_2 and S_2O_1 . The objective is to minimize the differential error between the two convolution seismograms, and achieve an optimization by cycling through the parameter space. Next we fix the RDP and DOBs of these two explosions using the derived optimized values and continue the process to include the next explosion. Thus, we have two additional source convolved seismograms: S_3O_1 and S_3O_2 to minimize the global error between these constructed differential seismograms. We continue the process until the last explosion is included. The investigation resulted RDP and DOB parameters consistent with those established by other investigators except for the September 3, 2017 explosion, which is caused by the influence of non-isotropic seismic sources, and which is a topic of current investigation.

Primary author: SAIKIA, Chandan (U.S. Air Force Technical Applications Center)

Presenter: SAIKIA, Chandan (U.S. Air Force Technical Applications Center)

Session Classification: T2.3 Seismoacoustic Sources in Theory and Practice

Track Classification: Theme 2. Events and Nuclear Test Sites