



ID:

Type: Poster

## and evolution of global Xe background between 2016 and 2018

Monitoring atmospheric radionuclides is essential for confirming an underground nuclear weapon test. Accordingly, it is crucial to the international monitoring system (IMS) of the Comprehensive Nuclear-Test-Ban Treaty (CTBT) to monitor radionuclides worldwide. Four of the xenon isotopes are of interest for CTBT verification;  $^{131m}\text{Xe}$ ,  $^{133m}\text{Xe}$ ,  $^{133}\text{Xe}$  and  $^{135}\text{Xe}$ . In a nuclear explosion, the four isotopes are produced in sufficient quantities and have half-life times long enough to allow an appreciable amount to travel and reach IMS radionuclide stations. Therefore, understanding the Xenon background at the IMS stations is important to CTBT verification process. In this work, a comparative study between the global radionuclide background in 2016, 2017 and 2018 is conducted to determine the evolution of Xe background for all the available IMS stations. The stations with pronounced changes in its Xe background were identified and the ratios between different Xe isotopes were calculated and analyzed.

**Primary author:** ELBAHRAWY, Mohammed Yehia Taha Ahmed (National Research Institute of Astronomy and Geophysics (NRIAG))

**Presenter:** ELBAHRAWY, Mohammed Yehia Taha Ahmed (National Research Institute of Astronomy and Geophysics (NRIAG))

**Track Classification:** Theme 2. Events and Nuclear Test Sites