



ID:

Type: **Poster**

## **New Detection Mediums for Atmospheric Radioxenon Measurements**

Several radioxenon isotopes ( $\text{Xe-131m}$ ,  $\text{Xe-133}$ ,  $\text{Xe-133m}$ ,  $\text{Xe-135}$ ) are characteristic byproducts of nuclear explosions, and the presence of these isotopes in specific ratios in the atmosphere acts as a clear tracer which allows the Comprehensive Nuclear Test Ban Treaty Organization (CTBTO) to verify the nuclear nature of a clandestine explosion. These isotopes can be discriminated from background and detected at extremely low concentrations ( $< 1 \text{ mBq/m}^3 \text{ air}$ ) via exploitation of their distinct beta-gamma coincidence decay signatures. At Oregon State University, we have recently developed three compact and relatively low-cost radioxenon detectors to improve reliability and maintainability of current radioxenon detection systems employed at the International Monitoring System. Our detectors utilize new detection mediums (Stilbene+SiPM,  $\text{SrI}_2$ +SiPM, co-planar CZT, and PIPS detectors) to measure xenon radioisotopes via beta-gamma coincidence technique. In this presentation, we will present the design of detectors and also discuss our preliminary experimental results using (1) Stilbene-CZT, (2) PIPS-CZT, and (3) PIPS- $\text{SrI}_2$  radioxenon detectors.

**Primary author:** FARSONI, Abi (Oregon State University)

**Presenter:** FARSONI, Abi (Oregon State University)

**Track Classification:** Theme 3. Verification Technologies and Technique Application