

Low Frequency Acoustical Propagation in a Shallow-Water Wedge

Tuesday, 20 June 2023 09:01 (1 minute)

In shallow-water environments long range propagation proceeds by repeated reflections from the surface and the bottom of the channel, as is the case for underwater sound of a wide spectral range, whose very low frequencies may propagate over large distances, without significant losses. In this paper, a 3-D benchmark model of the fluid wedge over an elastic bottom is applied to explain low frequency long range propagation in shallow water overlaying a sloping elastic-type seabed, such as a sedimentary rock exhibiting rigidity. The modeling approach is based on the modified method of generalized ray that furnishes a complete acoustic signal, thus comprised of contributions from all of the wave motions typical for the model (not only from the source signal and the regularly [specularly] reflected waves but also from the refracted [lateral] waves and the pseudo-Rayleigh and the Scholte interface waves), received in order of their arrivals at a large distance from the point source. When the source emits signals of a low frequency content, the contribution from the Scholte waves becomes dominant at large distances. Hence, low frequency long range propagation in a shallow-water wedge (coastal wedge) may indeed be governed by the Scholte waves.

E-mail

piotr.borejko@tuwien.ac.at

Promotional text

Low frequency long range acoustical propagation in a shallow-water wedge (coastal wedge) may indeed be governed by the Scholte waves.

Oral preference format

in-person

Primary author: BOREJKO, Piotr (TU Wien)

Presenter: BOREJKO, Piotr (TU Wien)

Session Classification: Lightning talks: P1.3, P1.4, P5.2

Track Classification: Theme 1. The Earth as a Complex System: T1.3 The Oceans and their Properties